Fen and Marsh Habitat Action Plan

1. Introduction

Fen and marsh vegetation is groundwater-fed permanently, seasonally or periodically waterlogged peat, peaty or mineral soils where grasses do not predominate. It also includes emergent vegetation or frequently inundated vegetation occurring over peat or mineral soils. It does not include areas of carr that are greater than 0.25 ha nor wet grassland (with the exception of purple moor grass, reed, or sweet-grass dominated vegetation). Lowland wet grassland has its own Action Plan within this BAP

The following relevant habitats were listed as UK BAP priority habitats and subsequently in Section 41 of the Natural Environment and Rural Communities (NERC) Act 2006: Lowland fens; Coastal and floodplain grazing marsh; Purple moor grass and rush pastures.

The UK is thought to host a large proportion of the fen surviving in the EU. As in other parts of Europe, fen vegetation has declined dramatically in the past century. Peatland habitats have been identified as major contributors to carbon storage and their degradation leads to the release of thousands of tonnes of CO_2 into the atmosphere every year.

Within the county fen and marsh, as with other wetland habitats, have undergone a serious decline in extent and quali

S11	Although bottle sedge often dominates this community in shallow water there can be other species such as soft rush, sometimes in reasonable amounts.
S12	*Typha latifolia* swamp (4.18ha) Common reedmace is always dominant, frequently with no other species present.
S13	*Typha angustifolium* swamp (0.56ha) This is dominated by lesser reedmace, which prefers more basic water around pools with silty substrate.
S14	*Sparganium erectum* swamp (1.33ha) This typical sub-community is normally species poor with the branched9.6 r 367.871 110.42 664.4rq161.18 608.26 367.87 39.6 reW*ı

	grazing.
WE27	*Epilobium hirsutum* weed community (1.36ha) Greater willowherb dominates this tall herb community on damp ground normally along riverbanks and in areas of ungrazed marsh.
Unknown	Dominants *Scirpus sylvaticus-Carex pseudocyperus* (0.76ha) On several sites, areas of swamp dominated by wood clubrush and cyperus sedge occur.
Total area	53.62ha

2.2

Puxton Marshes SSSI comprises a large area of unimproved marshy grassland with associated damp woodland and open water. The marsh is notable for the variety of its plants of which 110 species have been recorded. Most of the marshy grassland is dominated by tall fen with reed canary-grass (*Phalaris arundinacea)*, reed sweet-grass (*Glyceria maxima)*, common valerian (*Valeriana officinalis)*, great hairy willowherb (*Epilobium hirsutum)* and meadowsweet (*Filipendula ulmaria)*. Other less common species include marsh cinquefoil (*Potentilla palustris)*, skullcap (*Scutellaria galericulataini5.9(o)4(n)4(u)3(s-1(o)4(i)3)]TJ0T3T26.3 0 1 1205.07 2.5 Tm[())]TJ

common cotton-grass (*Eriophorum angustifolium)*, marsh cinquefoil and sphagnum mosses. The diverse flora and small pools help to support rare fauna including several regionally scarce species of Lepidoptera and Odonata. Water levels in The Bog appear to have lowered over time and actions have been taken to increase these such as tree removal. Investigations have also been made into artificially maintaining water levels from bore hole fed pumps.

3. Current factors affecting the habitat

Groundwater abstraction and/or field drainage lowering the water table so that many important fen and marsh sites are now drying out leading to changes in vegetation communit 5Tf1 erer1re6(re)Tfr23(5Tf0 TJET3)-169(l)-6(e)4(ad)]TJE

4.3 Survey, research and monitoring

Lee, J. A (1975). *The conservation of British inland salt marshes*. Biological Conservation, 8, pp 143-151.

Lee, J. A (1977). *The Vegetation of British Inland Salt Marshes*. Journal of Ecology, 65, pp. 673-698.

Liley, M (1999). *botanical survey.* Worcestershire Wildlife Trust.

Natural England (2015). *Hydrological functioning theme plan. Restoring the hydrology of Natura 2000 terrestrial wetlands.* Natural England.

Natural England & RSPB (2015). *Climate Change Adaptation Manual - Evidence to support nature conservation in a changing climate (NE546) 12 Lowland fens.* http://publications.naturalengland.org.uk/publication/5629923804839936.

McBride, A., Diack, I., Droy, N., Hamill, B., Jones, P., Schutten, J., Skinner, A and Street M (Eds) (2011). *The Fen Management Handbook*. Scottish Natural Heritage, Perth.

Wheeler, B. D., Gowing, D. J. G., Shaw, S. C., Mountford, J. O and Money R. P (2004). *Ecohydrological Guidelines for Lowland Wetland Plant Communities.* (Eds. Brooks A. W., Jose P. V., and Whiteman M. I). Environment Agency (Anglian Region).

Fens for the Future Partnership https://www.fensforthefuture.org.uk/the-partnership/